

## ON RIGHT PRINCIPALLY PROJECTIVE SKEW POWER SERIES RINGS

AHMAD MOUSSAVI

Departement of Mathematics,  
Tarbiat Modares University,  
P.O.Box. 14115-343, Tehran, Iran.  
`moussavi.a@modares.ac.ir`

Ali Ahmadi  
Departement of Mathematics, ACER,  
Tarbiat Modares University,  
P.O.Box. 14155-4838, Tehran, Iran  
`aliahmadi@modares.ac.ir`

ABSTRACT. In this paper we give a generalization of a result of J.A. Fraser and W.K. Nicholson [2]. Let  $R$  be a ring and  $\alpha$  be a weakly rigid automorphism of  $R$ . If  $R$  is skew power series Armendariz, then  $R[[x; \alpha]]$  is right principally projective if and only if  $R$  is right principally projective and any countable family of idempotents in  $R$  has a generalized join when all left semicentral idempotents are central.

### 1. INTRODUCTION

We want to construct polynomials over a (not necessarily commutative) ring  $R$  in one variable  $Y$  which needs not commute with elements of  $R$ . Further we want a unique representation for each non-zero polynomial of the form  $\sum_{i=0}^p r_i Y^i$ , where  $r_0, \dots, r_p \in R$  and  $r_p \neq 0$ , i.e. we want the polynomial ring to be a free left  $R$ -module with basis  $\{1, Y, Y^2, \dots\}$ . Degrees of polynomials, defined as

$$\deg \left( \sum_{i=0}^p r_i Y^i \right) := p,$$

(where  $r_p \neq 0$ ) shall be respected by multiplication, in the following sense:

$$\deg(fg) \leq \deg(f) + \deg(g).$$

This means in particular that for  $r \in R$  we want to have

$$Yr = \alpha(r)Y + \delta(r)$$

---

**2000 Mathematics Subject Classification:** 16S36, 16E50.

**keywords and phrases:** Baer rings; right p.p.-rings; weakly rigid automorphism; Armendariz rings; skew power series.

for some mappings  $\alpha, \delta : R \rightarrow R$ . Consequently  $\alpha$  and  $\delta$  need to have some special properties: By ring properties we conclude for  $r, r' \in R$

$$\alpha(r+r')Y + \delta(r+r') = Y(r+r') = Yr + Yr' = (\alpha(r) + \alpha(r'))Y + \delta(r) + \delta(r')$$

and

$$\begin{aligned} \alpha(rr')Y + \delta(rr') &= Y(rr') = (Yr)r' = (\alpha(r)Yr' + \delta(r)r') \\ &= \alpha(r)Yr' + \delta(r)r' = \alpha(r)(\alpha(r')Y + \delta(r')) + \delta(r)r' \\ &= \alpha(r)\alpha(r')Y + \alpha(r)\delta(r') + \delta(r)r' \end{aligned}$$

which, using module basis properties, implies

$$\alpha(r+r') = \alpha(r) + \alpha(r')$$

$$\alpha(rr') = \alpha(r)\alpha(r')$$

(i.e.  $\alpha$  is an endomorphism of  $R$ ) and

$$\delta(r+r') = \delta(r) + \delta(r')$$

and

$$\delta(rr') = \alpha(r)\delta(r') + \delta(r)r'$$

(i.e.  $\delta$  is an  $\alpha$ -derivation as defined below). It turns out, that these properties are not only necessary, but already sufficient for the existence of such a ring.

Throughout  $\alpha : R \rightarrow R$  is an automorphism and  $C(R)$  the center of  $R$ . We denote  $S = R[[x; \alpha]]$  the skew power series ring, whose elements are power series of the form  $\sum_{i=0}^{\infty} r_i x^i$  with coefficients  $r_i \in R$ , where the addition is defined as usual and the multiplication subject to the condition  $xb = \alpha(b)x$ , for any  $b \in R$ .

In [5] Rickart studied  $C^*$ -algebras with the property that every right annihilator of any element is generated by a projection (i.e.,  $p$  is a projection if  $p = p^2 = p^*$  where  $*$  is the involution on the algebra). A ring satisfying a generalization of Rickarts condition (i.e., every right annihilator of any element is generated (as a right ideal) by an idempotent) has a homological characterization as a right PP ring. A ring  $R$  is called a right (resp. left) PP ring if every principal right (resp. left) ideal is projective (equivalently, if the right (resp. left) annihilator of an element of  $R$  is generated (as a right (resp. left) ideal) by an idempotent of  $R$ ).  $R$  is called a PP ring (also called a Rickart ring if it is both right and left PP. There is a right p.p.-ring which is not right p.q.-Baer.

In 1974, Armendariz showed that a reduced ring  $R$  is Baer if and only if  $R[x]$  is Baer [1, Theorem B]. Armendariz also provided an example to show that the reduced condition is not superfluous.

A ring  $R$  is called *Armendariz* if whenever two polynomials  $f(x) = \sum_{i=0}^m a_i x^i$ ,  $g(x) = \sum_{j=0}^n b_j x^j \in R[x]$  satisfy  $f(x)g(x) = 0$  we have  $a_i b_j = 0$  for every  $i, j$ .

Fraser and Nicholson in [2] showed that  $R[[x]]$  is a reduced p.p.-ring if and only if  $R$  is a reduced p.p.-ring and any countable family of idempotents of  $R$  has a least upper bound in  $I(R)$ , the set of all idempotents.

Z. Liu in [4, Theorem 3], showed that: If  $R$  is a ring such that all left semicentral idempotents are central, then  $R[[x]]$  is right p.q.-Baer if and only if  $R$  is right p.q.-Baer and any countable family of idempotents in  $R$  has a generalized join in  $I(R)$ .

A monomorphism  $\alpha$  is said to be rigid if for each  $a \in R$ ,  $a\alpha(a) = 0$  implies that  $a = 0$ . According [3] a monomorphism  $\alpha$  is said to be weakly rigid if for each  $a, b \in R$ ,  $ab = 0$  implies that  $a\alpha(b) = \alpha(b)a = 0$ . By [3, Proposition 3], every rigid endomorphism is weakly rigid. By [3] there are examples of weakly rigid endomorphism which are not rigid.

2. SKEW POWER SERIES ARMENDARIZ RINGS

Motivated by results in Armendariz [1], we investigate a generalization of  $\alpha$ -rigid rings and introduce skew power series versions of the Armendariz rings:

**Definition 2.1.** For a ring  $R$  and an automorphism  $\alpha : R \rightarrow R$ , we say  $R$  is skew power series Armendariz, if for each  $f(x) = \sum_{i=0}^{\infty} a_i x^i$  and  $g(x) = \sum_{j=0}^{\infty} b_j x^j \in R[[x; \alpha]]$ ,  $f(x)g(x) = 0$  if and only if  $a_i b_j = 0$  for all  $i, j$ .

We first note that there is an example of a non reduced regular ring (hence p.p) that is neither right nor left p.q.-Baer.

**Lemma 2.2.** Let  $R$  be a weakly rigid ring. Then we have the following:  
 (i) If  $ab = 0$ , then  $aa^n(b) = \alpha^n(a)b = 0$  for each positive integer  $n$ .  
 (ii) If  $aa^k(b) = 0$  for some positive integer  $k$ , then  $ab = 0$ .

3. PRINCIPALLY PROJECTIVE SKEW POWER SERIES RINGS

In this section, we give a necessary and sufficient condition for some rings under which the ring  $R[[x; \alpha]]$  is right p.p.

**Proposition 3.1.** Let  $R$  be weakly rigid ring and  $S$  the skew power series ring  $R[[x; \alpha]]$ . Then the following statements are equivalent:  
 (1)  $R$  is skew power series Armendariz ;  
 (2)  $\varphi : rAnn_R(2^R) \rightarrow rAnn_S(2^S); A \rightarrow AS$  is bijective;  
 (3)  $\psi : \ell Ann_R(2^R) \rightarrow \ell Ann_S(2^S); B \rightarrow SB$  is bijective.

A ring is called *abelian* if every idempotent in it is central.

**Proposition 3.2.** Every skew power series Armendariz ring is abelian.  
**Definition 3.3.** (Z. Liu, [4]). Let  $\{e_0, e_1, \dots\}$  be a countable family of idempotents of  $R$ . We say  $\{e_0, e_1, \dots\}$  has a join in  $I(R)$  if there exists an idempotent  $e \in I(R)$  such that

1.  $e_i(1 - e) = 0$ , and
2. If  $f \in I(R)$  is such that  $e_i(1 - f) = 0$ , then  $e(1 - f) = 0$ .

**Theorem 3.4.** Let  $R$  be a weakly rigid skew power series Armendariz ring. Then  $R[[x; \alpha]]$  is right p.p. if and only if  $R$  is right p.p. and any countable family of idempotents in  $R$  has a join in  $I(R)$ .

**Corollary 3.5.**(J.A. Fraser, W.K. Nicholson [2]) Let  $R$  be a reduced power series ring. Then  $R[[x]]$  is right p.p. if and only if  $R$  is right p.p. and any countable family of idempotents in  $R$  has a join in  $I(R)$ .

## REFERENCES

- [1] E.P. Armendariz, A note on extensions of Baer and p.p-rings, *J. Austral. Math. Soc* **18** (1974) 470-473.
- [2] J.A. Fraser, W.K. Nicholson, Reduced PP-rings. *Math. Japonica* **34**(5) (1989) 715-725.
- [3] E. Hashemi and A. Moussavi , Polynomial Extensions of Baer and quasi-Baer rings, *Acta Math. Hungar.* **107** (3) (2005), 207224.
- [4] Z. Liu, A note on principally quasi-Baer rings, *Comm. Algebra* **30**(8) (2002) 3885-3890.
- [5] C.E. Rickart, Banach algebras with an adjoint operation. *Ann. of Math.* **47**(1946) 528-550.