

TRIANGULAR MATRIX REPRESENTATIONS OF RING EXTENSIONS

M. ROBAT SARPOOSHI AND E. HASHEMI

Shahrood University of Technology
 M.Sarpooshi@yahoo.com

ABSTRACT. We investigate the class of piecewise prime, *PWP*, rings which properly includes all piecewise domain. For a *PWP* ring we determine a large class of ring extensions which have a generalized triangular matrix representation for which the diagonal rings are prime. We investigate the quasi-Baer and related conditions on monoid rings $R[G]$. In particular we consider the transfer of the quasi-Baer and related conditions between R and a u.p.-monoid ring $R[G]$ of a u.p.-monoid G over R . We characterize the semi-central idempotents of various ring extensions of R in terms of the semi-central idempotents of R .

1. INTRODUCTION

All rings are associative and R denotes a ring with unity 1. By a ring extension we mean an covering of R which has the same unity as R . An idempotent $e \in R$ is left(right) *semi-central* in R if $exe = xe$ ($exe = ex$) for all $e \in R$. we use $S_l(R)$ and $S_r(R)$ for the sets of all left and right semi-central idempotents respectively. An idempotent e of R is *semi-central reduced* if $S_l(eRe) = \{0, e\}$. Note that $S_l(eRe) = \{0, R\}$ if and only if $S_r(eRe) = \{0, R\}$. A ring R is semi-central reduced if 1 is semi-central reduced. Also note that $S_r(R) \cap S_l(R) = B(R)$, where $B(R)$ is the set of all central idempotents R . A ring R has a *generalized triangular matrix representation* if there exists a ring isomorphism

$$\Theta : R \longrightarrow \begin{bmatrix} R_1 & R_{12} & \cdot & \cdot & \cdot & R_{1n} \\ 0 & R_2 & \cdot & \cdot & \cdot & R_{2n} \\ \cdot & 0 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & \cdot & \cdot & \cdot & 0 & R_n \end{bmatrix} \quad \text{where each diagonal ring, } R_i, \text{ is a}$$

ring with unity, R_{ij} is a left R_i -right R_j -bimodule for $i < j$, and the matrices obey the usual rules for matrix addition and multiplication. If each R_i is semi-central reduced then R has a *complete generalized triangular matrix representation* with triangulating dimension n . R is called *right principally quasi-Baer* (or simply *right p.q.-Baer*) if the right annihilator of a principal right ideal is

2000 Mathematics Subject Classification: 16S60;16D70;16S50.

keywords and phrases: Ring extension; Quasi-Baer ring; Semi-central idempotent; U.p.-monoid ring.

generated by an idempotent. An ordered set $\{b_1, \dots, b_n\}$ of nonzero distinct idempotents in R is called a set of *left triangulating idempotents* of R if all the following hold: (i) $1 = b_1 + \dots + b_n$; (ii) $b_1 \in S_l(R)$; and (iii) $b_{k+1} \in S_l(c_k R c_k)$ where $c_k = 1 - (b_1 + \dots + b_k)$ for $1 \leq k \leq n - 1$. A set $\{b_1, \dots, b_n\}$ of left triangulating idempotents is said to be *complete* if each b_i is semi-central reduced. Any complete set of primitive idempotents determines a complete set of left triangulating idempotents. R has a (complete) set of left triangulating idempotents if and only if R has a (complete) generalized triangular matrix representation. R has *triangulating dimension* n , written $Tdim(R) = n$ if R has a complete set of left triangulating idempotents with exactly n elements. R is semi-central reduced if and only if $Tdim(R) = 1$. Let B be a set of left triangulating idempotents of R and Γ a ring extension of R . (i) We say Γ is *B-triangularly linked* to R if whenever $b \in B$ and $0 \neq c \in S_l(b\Gamma b)$, then there exists $0 \neq c_0 \in S_l(bRb)$ such that $c_0\Gamma \subseteq c\Gamma$. (ii) we say Γ is *B-triangularly compatible* with R if B is a set of left triangulating idempotents of Γ . If Γ is B-triangularly linked to (resp. triangularly compatible with) R for every set B of left triangulating idempotents, then we say Γ is *triangularly linked to* (resp. *triangularly compatible* with) R . we call $b \in R$ a *triangulating idempotent* if b is in some set of left triangulating idempotents of R . For any subset X of a ring R , $r_R(X)$ (resp. $\ell_R(X)$) denotes the right (resp. left) annihilator of X in R .

2. MAIN RESULTS

Theorem 2.1. *Let $R[G]$ be the monoid ring of a u.p.-monoid G over a ring R . then we have the following: (i) R is right p.q.-Baer if and only if $R[G]$ is right p.q.-Baer. (ii) R is quasi-Baer if and only if $R[G]$ is quasi-Baer.*

Theorem 2.2. *let σ be a ring endomorphism of R and X a nonempty set of not necessarily commuting indeterminates. (i) Let $\Gamma = R[x; \sigma]$ or $R[[x; \sigma]]$. Then $e \in S_l(\Gamma)$ with $e_0 \in R$ the constant term of e if and only if $e_0 \in S_l(R)$, $ee_0 = e_0$, $e_0e = e$. (ii) Let $\Delta = R[X]$ or $R[[X]]$. Then $e \in S_l(\Delta)$ with $e_0 \in R$ the constant term of e if and only if $e_0 \in S_l(R)$, $ee_0 = e_0$, $e_0e = e$.*

Theorem 2.3. *Let R be a right p.q.-Baer ring. Then $Tdim(R) = n$ if and only if R has exactly n minimal prime ideals.*

Theorem 2.4. *Let 1_M denote the unity of $Mat_n(R)$ and $\bar{R} = R.1_M$. Then $e = (e_{ij}) \in S_l(Mat_n(R))$ if and only if $e_0 \in S_l(\bar{R})$ and $e_0 Mat_n(R) = e Mat_n(R)$, where $e_0 = e_{11}.1_M$.*

Theorem 2.5. *Let X be a nonempty set of not necessarily commuting indeterminates, σ an endomorphism of R , and G a monoid. The following extensions of R are triangularly linked to R and triangularly compatible with R , hence have the same triangulating dimension as R : (i) $R[G]$ where G is a u.p.-monoid and R is a right p.q.-Baer ring; (ii) $R[G]$ where G is a free monoid; (iii) $R[X]$; (iv) $R[[X]]$; (v) $R[x, x^{-1}]$; (vi) $R[[x, x^{-1}]]$; (vii) $R[x; \sigma]$ where σ is a ring endomorphism of R such that $\sigma(bR) \subseteq bR$ for every left triangulating idempotent $b \in R$; (viii) $R[[x; \sigma]]$ where σ is a ring endomorphism of R such that $\sigma(bR) \subseteq bR$ for every left triangulating idempotent $b \in R$; (ix) $Mat_n(R)$.*

Theorem 2.6. *Let R be a quasi-Baer ring with a complete set of left triangulating idempotents $B = \{b_1, \dots, b_n\}$. If Γ is any of the following ring extensions of R then Γ is a quasi-Baer ring with B determining a complete generalized triangular matrix representation for Γ in which each diagonal ring R_i is a prime ring: (i) $R[G]$ where G is a u.p.-monoid; (ii) $R[X]$ where X is a nonempty set of not necessarily commuting indeterminates; (iii) $R[[X]]$ where X is a nonempty set of not necessarily commuting indeterminates; (iv) $R[x, x^{-1}]$; (v) $R[[x, x^{-1}]]$; (vi) $R[x; \alpha]$ where α is a ring automorphism such that $\alpha(bR) \subseteq bR$ for all $b \in R$; (vii) $R[[x; \alpha]]$ where α is a ring automorphism such that $\alpha(bR) \subseteq bR$ for all $b \in R$; (viii) $T_n(R)$; (ix) $Mat_n(R)$.*

Theorem 2.7. *For a ring R with a complete set of triangulating idempotents, the following conditions are equivalent: (i) R is right p.q.-Baer; (ii) R is left p.q.-Baer; (iii) R is quasi-Baer; (iv) R is a pwp ring.*

Lemma 2.8. *Let $R = F[G]$ be a semi-prime group algebra over a field F . Then R is quasi-Baer if and only if each annihilator ideal is finitely generated.*

Lemma 2.9. *Let $\{b_1, \dots, b_n\}$ be a set of left triangulating idempotents of R . Then $Tdim(R) = \sum_{i=1}^n Tdim(b_i R b_i)$. In particular $n \leq Tdim(R)$.*

Lemma 2.10. *If R satisfies any of the following conditions then R has a complete triangular matrix representations: (i) R has a complete set of primitive idempotents; (ii) R has no infinite set of orthogonal idempotents; (iii) R has D.C.C on (idempotent generated, principal, or finitely generated) ideals; (iv) R has D.C.C on (idempotent generated, principal, or finitely generated) ideals; (v) R has A.C.C on (idempotent generated, principal, or finitely generated) ideals; (vi) R has A.C.C on (idempotent generated, principal, or finitely generated) ideals; (vii) R has either A.C.C or D.C.C on right annihilators; (viii) R is a semi-primary ring.*

Lemma 2.11. *The following conditions are equivalent: (i) R has a complete set of left triangulating idempotents; (ii) $\{bR \mid b \in S_l(R)\}$ is a finite set; (iii) $\{bR \mid b \in S_l(R)\}$ satisfies A.C.C and D.C.C; (iv) $\{bR \mid b \in S_l(R)\}$ and $\{Rc \mid c \in S_r(R)\}$ satisfies A.C.C; (v) $\{bR \mid b \in S_l(R)\}$ and $\{Rc \mid c \in S_r(R)\}$ satisfies D.C.C; (vi) $\{bR \mid b \in S_l(R)\}$ and $\{cR \mid c \in S_r(R)\}$ satisfies D.C.C; (vii) $\{bR \mid b \in S_l(R)\}$ is a finite set; (viii) R has a complete set of right triangulating idempotents; (ix) R has a complete generalized triangular matrix representation;*

Lemma 2.12. *If $\{e_1, \dots, e_n\}$ is a complete set of primitive idempotents R , then $Tdim(R) \leq n$.*

Lemma 2.13. *A ring R is prime if and only if is quasi-Baer and semi-central reduced.*

Lemma 2.14. *A ring R is right p.q.-Baer if and only if whenever I is a principal ideal of R there exists $e \in S_l(R)$ such that $I \subseteq Re$ and $r(I) \cap Re = (1 - e)Re$.*

Lemma 2.15. *The following conditions are equivalent: (i) R is a prime ring; (ii) R is a semi-central reduced quasi-Baer ring; (iii) R is a semi-central reduced p.q.-Baer ring; (iv) R is a semi-central reduced right p.q.-Baer ring.*

Corollary 2.16. *Any PWD (piecewise domain) is a quasi-Baer ring.*

Corollary 2.17. *If ring R is noetherian or artinian ring then R has a triangular matrix representation.*

Corollary 2.18. *Let Γ be a ring extension of R such that $\Gamma I \subseteq I\Gamma$ for all ideal I of R and whenever $b \in S_l(\Gamma)$, then there exists $b_0 \in R$ such that $b_0b = b$ and $bb_0 = b_0$. If Γ is prime (resp. left p.q.-Baer) then R is prime (resp. left p.q.-Baer).*

Corollary 2.19. *If $Tdim(R) \lesssim \infty$, then if R is prime then R is right p.q.-Baer.*

Corollary 2.20. *A ring R is prime with $Tdim(R) = m$ if and only if $T_n(R)$ is prime with $Tdim(T_n(R)) = mn$.*

Corollary 2.21. *The PWP property is a Morita invariant.*

Corollary 2.22. *Let A and B be semi-central reduced rings. If $T_m(A) \cong T_n(B)$ for some m and n , then $m = n$ and $A \cong B$*

Corollary 2.23. *The right p.q.-Baer condition is a Morita invariant property.*

REFERENCES

- [1] G. F. Birkenmeier, J. K. Park *Triangular matrix representations of ring extensions*, J. Algebra **265** (2003) 457-477.
- [2] G. F. Birkenmeier, H. E. Heatherly, J. Y. Kim, J. K. Park *Triangular matrix representations*, J. Algebra **230** (2000) 558-595.
- [3] G. F. Birkenmeier, J. Y. Kim, J. K. Park, *Principally quasi-Baer rings*, Comm. Algebra **29** (2001) 639-660.
- [4] N. Groenewald *A note on extensions of Baer and p.p.-rings*, Publ. Linstiute Math. **34** (1983), 71-72.